

# Bill harnesses on nestling Tufted Puffins influence adult provisioning behavior

Carina Gjerdrum,<sup>1,5</sup> Gregor M. Yanega,<sup>2</sup> and Douglas F. Bertram<sup>1,3,4</sup>

<sup>1</sup>Department of Zoology, University of British Columbia, Vancouver, BC, Canada; <sup>2</sup>Department of Biology, University of Victoria, Victoria, BC, Canada; <sup>3</sup>Department of Zoology, University of Guelph, Guelph, ON, Canada; <sup>4</sup>Department of Zoology, University of Toronto, Toronto, ON, Canada; <sup>5</sup>Department of Zoology, University of Alberta, Edmonton, AB, Canada

Received 9 September 2005; accepted 17 February 2006

**ABSTRACT.** For burrow-nesting seabirds, investigators have examined nestling diet by attaching harnesses to the bills of nestlings to intercept food delivered by the parent. To determine whether this method provides an unbiased estimate of nestling diet, we evaluated its effect on the provisioning behavior of Tufted Puffins (*Lophochochus auritus*) nesting on Triangle Island, British Columbia. Adults delivering food to nestlings with bill harnesses always hesitated before entering a burrow with food, increasing their susceptibility to kleptoparasitism and did not always leave the food intended for the nestling. These responses by adult puffins could lead to underestimates of energy intake rates of nestlings and unreliable comparisons with other species if prey left by adults in nest burrows were the only source of data. We also compared estimates of the species, number, and size of prey delivered by adult puffins as determined by direct observation from blinds to samples of prey collected directly from nest burrows and found that the two sampling techniques produced similar results. However, identifying rare prey species and gathering precise information about prey length, mass, and condition require collection of prey, and we recommend

éndonse a kleptoparasitismo por parte de gaviotas. Además no siempre le dejaron comida a los pichones. La respuesta de los adultos, puede llevar a subestimar las necesidades energéticas de los pichones y hacer comparaciones poco confiables con otras especies si se utiliza únicamente como datos, las presas dejadas en las guaridas por los adultos. También comparamos las especies utilizadas para alimentar a los polluelos, número de presas y su tamaño comparando observaciones directas de lo que se llevaba a los polluelos con muestras de presas recolectadas directamente de las guaridas y encontramos que los dos métodos produjeron resultados similares. Sin embargo, identificar especies de presas raras y obtener información precisa sobre la especie, longitud, peso y condición de esta, requieren el examinar las misma. Recomendamos utilizar una combinación de

both methods to obtain reliable estimates of nestling diet, fish, *Lophochochus auritus*, ligature, sampling prey, seabird

least disturbance, but still provide reliable, unbiased, and repeatable estimates. Because human disturbance of adult Tufted Puffins (*Fratercula cirrhata*) can cause them to abandon nests (Pierce and Simons 1986), several investigators have used harnesses attached to the bills of nestlings to intercept prey delivered by parents (Hatch 1984, Baird 1986, 1990, 1991, Kitaysky 1996).

Tufted Puffins breed on islands along the Pacific Rim from California to Hokkaido, Japan, and are most abundant in British Columbia, Alaska, and in the Sea of Okhotsk (Piatt and Kitaysky 2002). Females lay a single egg in a burrow, and both parents feed the nestling several times a day with prey carried crosswise in the bill. Nestling diets are dominated by fish, although there are regional and interannual differences in prey size and species composition (Vermeer 1979, Wehle 1983, Hatch 1984, Baird 1990, 1991, Hatch and Sanger 1992, Piatt et al. 1997, Kitaysky and Golubova 2000, Gjerdrum 2004).

To reliably estimate nestling diet, the effect that a harnessed chick has on parental provisioning behavior needs to be evaluated. Our specific objectives were to (1) describe any behavioral differences between adult Tufted Puffins delivering food to nestlings with and without bill harnesses, and (2) compare visual estimates of bill loads (prey in the bill of adults being delivered to nestlings) to samples collected from burrows.

## METHODS

We studied puffins in the Puffin Rock sub-colony on Triangle Island, British Columbia, Canada (50°52'N, 129°05'W), during the breeding seasons of 1999 and 2000 when fledging success was unusually high (Gjerdrum et al. 2003). In both years, as part of a larger study on nestling growth and parental provisioning behavior, approximately 100 burrows were marked with a flag that could be read from an observation blind located 50–100 m from the burrows. If the nestling could not be reached from the entrance of the burrow, an access hole was dug and covered with a cedar shingle, dirt, and grass.

On 23 July, 29 July, and 2 August 1999, harnesses made of twist ties and cotton string were attached to the bills of 10 nestlings, and the same nestlings were manipulated each time. Twist ties were securely fastened around the bill, distal to the nares, and held in place with the

string tied around the back of the head (Baird 1986). Harnesses remained on the nestlings for 4 h and prevented them from swallowing food, but not from vocalizing. On each sampling date, harnesses fell off 2 of the 10 nestlings before the end of the observation period and, as a result, our sample size of harnessed nestlings for each sampling period was 8. Harnessed nestlings occupied burrows located among an additional 62 active burrows.

We monitored all burrows from an observation blind, and counted feeding visits and recorded parental behavior from 06:00 to 10:00. After the observation period, harnesses were removed and prey were identified, weighed, measured, and then returned to the nestling. Prey mass was measured to the nearest 0.1 g using a spring scale (Pesola, Baar, Switzerland) and fish length was measured from the caudal peduncle to the end of the snout (standard length) using dial calipers (±0.1 mm).

In 2000, we compared visual estimates of bill loads to samples collected from burrows. Between 13 July and 17 August, we estimated bill load size and composition from an observation blind using 8×30 binoculars. When possible, we estimated the number, size, and species of prey in food deliveries during 13 4-h observation periods. Lengths of fish were estimated based on their relationship to the size of the puffin bill. Two distinct size-classes of sand lance (*Ammodytes americanus*; juvenile <105 cm < adult) and one size-class of rockfish (*Sebastes* spp.; juvenile <60 cm) were identified. The mass of bill loads was estimated using species-specific length-to-mass relationships determined from collected samples (Gjerdrum 2001). When fish species could not be determined, we assigned weight based on an average mass for all fish collected in that size class.

To compare visual estimates of bill load size and composition to prey samples collected from burrows, we attached bill harnesses to chicks on 13 July (n = 10), 20 July (n = 16), 27 July (n = 16), and 4 August (n = 9). Harnesses were left on for 24 h to maximize the probability that parents would leave food intended for the chicks. The same nestlings were harnessed on each date. We identified and measured prey found in the burrows and subsequently fed them to the nestlings. Prey either dropped by puffins being chased by Gulls or left in burrows (when nestlings were pulled from burrows to be

measured and weighed) between 13 July and 17 August were also measured ( $n = 29$ ).

We used  $\chi^2$  analyses to determine if the relative proportion of prey species delivered by parents differed between samples estimated visually during delivery and those that were intercepted. We present means  $\pm 1$  SD. SYSTAT 8.0 (SPSS Inc. 1998) was used for all analyses.

## RESULTS

During the 4-h observation period, adults delivered food during 25%, 67%, and 71% of all feeding visits on the three sampling dates, respectively (Table 1). All adults hesitated for at least 30 s at the burrow entrance before entering. For the failed feeding attempts, adults quickly left burrows still carrying food, and three food loads were subsequently kleptoparasitized by Glaucous-winged Gulls (*Larus glaucescens*). On 2 August, one food load was delivered to the nestling only after two previous attempts (i.e., the parent came out of the burrow twice still carrying the food), and another was only a partial bill load because the parent flew away with some of the prey items. In contrast, we recorded 48–50 feeding visits to control nestlings on each of the sampling dates and only one adult failed to leave food for the nestling; the food-laden adult was chased by a Glaucous-winged Gull before entering its burrow.

We obtained 11 complete bill loads from the 21 feeding attempts (52% success) in 1999

Table 1. Behavior of Tufted Puffin parents feeding nestlings (

Table 2. Bill load composition of Tufted Puffins expressed as percentages (numerical abundance and wet mass for major prey species) of all items delivered.

Prey	Bill harness		Observation	
	% abundance	% mass	% abundance	% mass
Sand lance (0) <sup>1</sup>	8.2	7.8	8.6	7.7
Sand lance (1 ) <sup>2</sup>	4.3	15.9	4.6	16.8
Rockfish <sup>3</sup>	84.0	65.3	84.8	66.9
Squid <sup>4</sup>	1.2	3.3	0.7	0.2
Other invertebrates <sup>5</sup>	1.6	0.5	0.6	0.1
Other fish <sup>6</sup>	0.8	7.3	0.8	6.6
Number of samples	69		341	
Total prey items	257		1450	
Total mass (g) <sup>7</sup>	519.6		2938.8	

Data were obtained using bill harnesses and observation of provisioning adults from 13 July to 17 August 2000.

<sup>1</sup> *Ammodytes americanus* <105 mm fork length (Hatch and Sanger 1992).

<sup>2</sup> *Ammodytes americanus* >105 mm fork length (Hatch and Sanger 1992).

<sup>3</sup> *Sebastes* spp.

<sup>4</sup> *Loligo* spp.

<sup>5</sup> Includes octopus and larval fish.

<sup>6</sup> Includes Sablefish (*Anchoa mitchilli*) and any unidentified fish species.

<sup>7</sup> Bill load mass for observational samples was estimated using species-specific length-to-mass relationships (Gjerdrum 2001).

The delayed entry of adults into burrows with harnessed nestlings, and the rapid departures of adults still carrying food, suggest that parent-offspring communication facilitates successful feeding. Because nestlings with harnesses could vocalize, parents were likely reacting to calls that may have signaled alarm or distress. An audio recording of one harnessed nestling revealed persistent calling for the duration of a 60-min tape. By comparison, recordings revealed that nestlings without harnesses vocalized only when parents arrived with food (CG, unpubl. data). In general, parent-offspring communication in Tufted Puffins is not well understood. The function of chick vocalizations, variation among calls, and the environmental effects on communication require further study.

Previous investigators have assumed that placing bill harnesses or hoods on nestlings did not affect the behavior of parents (Baird 1990, 1991, Bertram et al. 1991, Bertram and Kaiser 1993, Kitaysky 1996). We have shown that food intended for nestlings is not always left by adults, and that the use of harnesses may increase the success of kleptoparasites. This bias could lead to underestimates of energy intake rates and unreliable comparisons among species or age

classes. If parents provision based on nestling nutritional requirements (Hamer and Hill 1994, Bertram et al. 1996, Harding et al. 2002, Gjerdrum 2004) and chicks vocalize to signal their nutritional needs to parents (Harris 1981), bill harnesses may also influence what parents bring to nestlings on subsequent feeding visits. Because we manipulated the same nestlings on each sampling date, parents may have habituated to the disturbance and been more willing to leave food for nestlings in the later sampling periods. Habituation to the method could introduce bias in analyses of the possible effects of season or nestling age on diet.

Visually estimating bill load size and composition causes the least disturbance and data we collected by observing adult puffins from blinds did not differ from that based on prey collected from burrows. Similarly, Rodway and Montevecchi (1996) found that visual observations of prey delivered by adult Atlantic Puffins (*Fregata aethiops*) provided reliable estimates of prey species composition. In 2000, rockfish and sand lance dominated the diet of Tufted Puffins on Triangle Island and could easily be distinguished using binoculars. However, a more diverse diet may have made it difficult to get reliable estimates

using this technique. For example, rare prey species such as herring (*Clupea pallasii*) or greenling (*Pachycottus* spp.) could be misidentified in bill loads with multiple species because these species are difficult to recognize from a distance. Squid, larval fish, and large euphausiids could be visually identified in the bill of puffins in this study, but smaller prey items make it more difficult to count individuals. Although larger sample sizes can be obtained by direct observation than by more disruptive and time-consuming interception methods, information on prey length and mass, prey condition, and identification of rare species require the collection of prey (Rodway and Montevecchi 1996).

Despite the effect bill harnesses had on the behavior of feeding parents, the technique is reliable for sampling species composition and for comparisons among years or colonies or areas of the same colony. This technique should also provide adequate sample sizes because the number of samples obtained per harnessed nestling is relatively high, depending on the length of time nestlings are left harnessed. However, several adult Tufted Puffins in Alaska stopped feeding their harnessed nestlings, and the technique led to high rates of nestling mortality (Hatch 1984). Because we found a high degree of similarity between visual and harness samples, we suggest using observations to estimate nestling diet, and supplementing the information with more invasive methods. This will minimize the potentially negative effects of harnesses, especially in years when food availability is low, on species of conservation concern like the Tufted Puffin.

#### ACKNOWLEDGMENTS

We thank J. Ryder for his assistance in the field and C. Cassady St. Clair for her initial descriptions and illustrations of the bill harness. Additional thanks to Ecological Reserves (B. C. Parks) for allowing access to Triangle Island (Anne Vallee Ecological Reserve). The Canadian Coast Guard provided ship and helicopter support with additional service provided by West Coast Helicopters. Logistical support for the field seasons were provided by the Nestucca Trust Fund Grant to D. F. Bertram, Natural Sciences and Engineering Research Council of Canada (NSERC) grants to F. Cooke, the Centre for Wildlife Ecology at Simon Fraser University (SFU), and the Canadian Wildlife Service. An SFU Graduate Fellowship and an NSERC postgraduate scholarship to CG provided additional financial support. Reviews by G. Ritchison, C. Cassady, St. Clair, and S. Hatch greatly improved the manuscript.

#### LITERATURE CITED

- AINLEY, D. G., G. BALLARD, K. J. BARTON, B. J. KARL, G. H. RAU, C. A. RIBIC

of Rhinoceros Auklets in Alaska. In: *Marine birds: their feeding ecology and commercial fisheries relationships* (D. N. Nettleship, G. A. Sanger, and P. F. Springer, eds.), pp. 106–115. Canadian Wildlife Service Special Publication, Ottawa, ON, Canada.

———, AND G. A. SANGER. 1992. Puffins as samplers of juvenile pollock and other forage fish in the Gulf of Alaska. *Marine Ecology Progress Series* 80: 1–14.

HEDD, A., J. R. R