

A

FLEMMING R. MERKEL,^{1,2}

KNUD FALK,

SARAH E. JAMIESON,


, 3900,

, 3900,

B



, B E3B 6E1, C

Ab ac



Ⓚ

A borealis, Somateria mollissima



¹ E- @ @ . g
²

F
 Su Ae a

O
 G (F)
 G
 C N G (B) 2004)
 N
 (M 2002). 507 56,735
 4,000 2,486
 D 32
 (579 ± 07 (E)
 H F G H D C
 (M 2002). 30
 (M 2004). 2002, 30
 O 3 M

M

C ec , X- a a d D ec
 A N M 3
 (2000 2002) N
 762 B
 ()
 (86%) (4%)
 A (0%)
 (: 5.2 6.0)
 (: 2.7 3.9)
 G (F 2006),
 (592)
 N A (C 2E)
 (35 × 40)
 N (470)
 40
 F (B 993, M E 999)

(B 963).
 H
 () ()
 0. (ING)

• • • • • • • • • •

- breeding dark-bellied Brent geese *B. hutchinsoni* in the high Arctic. *Journal of Avian Biology* 26:105–113.
- Falk, K., F. R. Merkel, K. Kampp, and S. E. Jamieson. 2006. Embedded lead shot and infliction rates of common and king eiders wintering in Southwest Greenland. *Wildlife Biology* 12:257–265.
- Fournier, M. A., and J. E. Hines. 1994. Effects of starvation on muscle and organ mass of king eiders and the ecological and management implications. *Wildfowl* 45:188–197.
- Frich, A. S., and K. Falk. 1997. Jagtindsats og ederfuglefangst ved Nuuk. Greenland Institute of Natural Resources, Technical Report No. 5, Nuuk, Greenland. [In Danish. Nuuk, Gree(Grie-eeenlans7soh-487.335(implicat)10.9(ei(K.)(RSomatvp(nd)-g7P5h(K.tidi95e)-239.2(E.)-235.6(Jamieson.