

exists when counting adult Piping Plovers outside a census window as our observations indicate not all birds retain the same mate or beach in a nesting season.

ACKNOWLEDGMENTS

The Canadian Wildlife Service funded the Piping Plover banding and monitoring program during the 2000 season. The Nova Scotia Department of Natural Resources provided in-kind and logistical support for the 2000 seasonal effort. Andrew Boyne and Kevin Davidson provided useful comments on earlier drafts of the manuscript.

REFERENCES CITED

- BATCHELOR, J. S., P. H. TRENKLE, G. DICKEY, R. WILSON, and D. SAUNDERS. 1994. Nova Scotia Piping Plover Atlas. Unpublished Nova Scotia Department of Natural Resources report. 86 pages.
- LITVIN, A. T. and C. H. CUMMINGS. (marked * in text)
- BATCHELOR, J. S., P. H. TRENKLE, G. DICKEY, R. WILSON, and D. SAUNDERS. 1994. Nova Scotia Piping Plover Atlas. Unpublished Nova Scotia Department of Natural Resources report. 86 pages.
- LITVIN, A. T. and C. H. CUMMINGS. (marked * in text)
- BATCHELOR, J. S., A. M. COOPER and B. LAURIE. 1997. Piping Plovers produce two broods. *Wilson Bulletin* 109: 337-339.
- CUMMINGS, W. E. 1977. Breeding biology of the piping plover in southern Nova Scotia. M.Sc. thesis, Dalhousie University, Halifax. 115 pages.
- CUMMINGS, W. E. 1982. Biology and behavior of breeding Piping Plovers. *Wilson Bulletin* 94: 531-545.
- HAY, S. M. 1987. The population biology and life history patterns of the piping plover. Ph.D. dissertation, University of North Dakota, Grand Forks. 121 pages.
- HAY, S. M. 1992. Piping Plover. Pages 1-18. *The Birds of North America* (2). A. Poole, P. Stettenheim,

and F. Gill. The Academy of Natural Sciences, Philadelphia; The American Ornithologists' Union, Washington, D.C.

HAY, S. M., and

on several occasions feeding on Tailed Frog (*Ascophryne*) larvae.

tant prey for larval Pacific Giant Salamanders (*Desmognathus siskiyou*) and Cope's salamanders (*Ambystoma macrodactylum*) and Red-legged Frogs (*Rana aurora*) (Bury 1968; Jones