RESEARCH ARTICLE

Are there synergistic or antagonistic effects of multiple maternally derived egg components on offspring phenotype?

Roxana Torres

sex communal cages with visual and acoustical contact with birds of the opposite sex, until assays to estimate breeding success were performed. The sex of surviving nestlings was determined at 60–

significant maternal treatment×sex interaction ($F_{1,52}$=6.09, P=0.017; egg mass, P<0.001; Table S1), so mass was analysed for each sex separately. For females, there was no main effect of MAb treatment ($F_{1,13}$=1.05, P>0.30), yolk testosterone treatment ($F_{1,13}$=0.69, P>0.40) or their interaction ($F_{1,13}$=0.65, P>0.40) on day 30 body mass (Fig. 2A). For males, there was no effect of yolk testosterone treatment or the interaction term (P>0.20) but there was a marginally significant effect of MAb treatment ($F_{1,12}$=4.22, P=0.06; Fig. 2): male chicks from MAb eggs were heavier than those from control eggs (13.6±0.2 versus 13.1±0.2 g). In a reduced model, excluding yolk testosterone treatment, there was a significant MAb treatment×sex interaction ($F_{1,56}$=8.49, P<0.01): female offspring from MAb eggs had a similar 30 day mass to controls (t_{56}=0.86, P=0.39), but male offspring from MAb eggs had a higher body mass than controls (t_{56}=2.60, P=0.012; Fig. 2B). For tarsus length at day 30 and body mass at maturity (day 82; Table 3), there was no effect of maternal treatment, egg treatment, sex, or any interaction (P>0.09 in all cases; Table S1).

Overall hatching success was independent of MAb treatment ($F_{1,254}$=1.62, P>0.20), yolk testosterone treatment ($F_{1,254}$=1.16, P>0.25) and their interaction ($F_{1,254}$=0.26, P>0.60; egg mass, P>0.30; Table 3). Similarly, fledging success was independent of MAb treatment ($F_{1,254}$=1.06, P>0.30), yolk testosterone treatment ($F_{1,254}$=0.51, P>0.45) and their interaction ($F_{1,254}$=0.44, P>0.50; egg mass, P=0.029; Table 3). Mean brood size at fledging did not vary among treatments (χ^2=1.05, d.f.=3, P>0.75; Table 3).

Effects of MAb and yolk testosterone treatment on offspring immune response

For the PHA response at fledging (day 26), there was a marginally significant main effect of yolk testosterone treatment ($F_{1,52}$=3.25, P=0.08; Fig. 3A,B) but the PHA response was independent of other main effects and interactions (P>0.20 in all cases; body mass, P=0.09; Table S2). In contrast, for the PHA response at maturity (day 83), there was a main effect of MAb treatment ($F_{1,42}$=7.19, P=0.010; Fig. 3C,D) but there were no other significant main effects or interactions (P>0.25 in all cases; body mass, P>0.55; Table S2). In both cases, the PHA response was higher in testosterone-treated eggs (Fig. 3B) or eggs from MAb-treated mothers (Fig. 3D) compared with controls.

The $\log_{10}$ LPS Ab titres in offspring at fledging (day 26) and at maturity (day 83) were independent of all main effects and interactions (P>0.15 in all cases; Table S3).

Effect of MAb and yolk testosterone treatment on adult male (sons) courtship behaviour

Courtship trials were conducted on n=49 male offspring at sexual maturity (day 90); on the first trial, n=30 males showed some courtship behaviour (i.e. 'invitation') and n=19 showed no courtship behaviour. For this first mating trail, there was no effect of MAb treatment ($F_{1,18}$=0.68, P>0.40) or a treatment interaction ($F_{1,18}$=0.18, P>0.60). However, a greater proportion of males from testosterone-treated eggs showed courtship behaviour (81.0%, n=21) compared with males from control eggs (46.4%, n=28; $F_{1,18}$=5.46, P=0.031). Of the 19 males tested in a second trial, 8 performed some courtship behaviour. Data were therefore pooled for these 8 males from the second trial and the 30 first trials for subsequent analysis of specific courtship behaviour (n=38 males; Table 4).

There was no overall difference in time to first copulation among the four combined MAb/yolk testosterone treatments (χ^2=4.83, d.f.=3, P=0.18; Table 4). However, after pooling data by egg treatment, time to first copulation was significantly shorter in testosterone-treated eggs compared with control eggs (Z=2.09, P=0.037; Fig. 4A). Conversely, after pooling data by MAb treatment, there was no difference in time to first copulation (P>0.90). For the number of successful mounts per 15 min there was a significant overall treatment effect (χ^2=8.15, d.f.=3, P=0.043). Males from control eggs had a higher number of successful mounts compared with males from testosterone-treated eggs (Z=2.06, P=0.039; Fig. 4B). However, the number of successful mounts was independent of MAb treatment (P>0.09). Male bill wiping behaviour and following behaviour were independent of combined treatments (P>0.30 in both cases; Table 4) and when MAb and yolk testosterone treatment were analysed separately (P>0.05 in all cases).

Effect of MAb and yolk testosterone treatment on adult female (daughters) reproduction

Among females surviving to sexual maturity (n=45) there was no effect of MAb, yolk testosterone treatment or their interaction on adult body mass ($F_{3,44}$=0.86, P>0.40) or tarsus length ($F_{3,35}$=1.06, P>0.30) at pairing.

Breeding propensity (percentage of females laying 1 egg) was high for all treatments: >80% (Table 5). There was no effect of MAb, yolk testosterone treatment or their interaction on the latency between pairing and laying of the first egg, clutch size or egg mass (P>0.12 in all cases; Table 5). Hatching success was independent of



MAb treatment ($F_{1,189}$=0.41, P>0.50), yolk testosterone treatment ($F_{1,189}$=1.83, P>0.15) and their interaction ($F_{1,189}$=0.12, P>0.70; Table 5). Similarly, fledging success was independent of MAb treatment ($F_{1,189}$=0.13, P>0.70), yolk testosterone treatment ($F_{1,189}$=0.01, P>0.90) and their interaction ($F_{1,189}$=0.73, P>0.35; Table 5).

Mean brood size at fledging did not vary among treatments (

Numerous studies have shown positive relationships between circulating antibody concentrations in mothers (but not fathers) and antibody concentrations in eggs or offspring (Gasparini et al., 2002; Grindstaff, 2008; Kowalczyk et al., 1985). In chickens, a secondary immune response to *E. coli* LPS peaked ~28 days after the first immunisation (Sunwoo et al., 1996); hence, by the time our females laid their first egg (≥21 days after the first injection), most of them should have been near the peak response phase to the immune challenge. Similarly, although we did not measure yolk testosterone, the dose of testosterone we injected corresponds to the difference in testosterone+dihydrotestosterone measured in yolks of eggs from females paired with attractive males versus unattractive males and thus mimics natural variation (Gil et al., 1999; von Engelhardt et al., 2006). This method of manipulating yolk testosterone has been widely used in zebra finches (Rutkowska et al., 2005; Tobler and Sandell, 2007) and other small passerines (Ruuskanen et al., 2009; Ruuskanen and Laaksonen, 2010; Tschirren et al., 2005) in studies of effects of maternal testosterone on offspring phenotype.

We acknowledge some potential limitations of our experimental design. Firstly, because of logistic constraints (~80 breeding pairs initially), we did not use a cross-fostering design, so eggs were laid and chicks reared by the same mother. However, we did use a

manipulation of maternal IgG had no effect on offspring growth in other studies (Burness et al., 2018; Grindstaff et al., 2006). Martyka et al. (2011) reported a sex-specific effect of maternal immunisation with sheep red blood cell antigen, but in the opposite direction to our result: daughters, but not sons, of immunised mothers were heavier and had longer tarsi (although chicks were measured at 12 days in their study, only half-way through the nestling growth period).

In our study, there was a positive effect of yolk testosterone on offspring immune response to PHA at 26 days of age, but at 82 days of age, MAb had a positive effect on offspring immune response to PHA. Previous studies have produced similarly mixed results; for example, Martyka et al. (2018) reported short-term immune-enhancing effects of MAb on nestling immune function, Merrill and Grindstaff (2014) reported immune-suppressing effects and Burness et al. (2018) found no effects of MAb on offspring immune function. Reid et al. (2006) reported long-term immune enhancement in offspring of vaccinated mothers up to 1 year after maternal treatment in free-living song sparrows, Melospiza melodia, but Addison et al. (2010) found no long-term effects of MAb in Japanese quail (Coturnix c. japonica). Most previous studies involving manipulation of yolk testosterone have found no or inconsistent effects on offspring immune function (reviewed in von Engelhardt and Groothuis, 2011; and see below).

The most compelling result we found was an effect of yolk testosterone on male courtship behaviour at sexual maturity, where a greater proportion of yolk testosterone-treated males engaged in courtship (81% versus 46%), and yolk testosterone-treated males had shorter times to first mounting and fewer successful mounts. Although the last two results might seem counter-intuitive, better males which mount females more quickly might need fewer attempts for functionally successful copulation (we defined successful mounts only as mounts with cloacal contact as we could not determine whether successful sperm transfer occurred). This would be consistent with positive, organisational effects of yolk androgens on dominance, competition and male sexual traits (Eising et al., 2006; Partecke and Schwabl, 2008; Schwabl et al., 2013; Strasser and Schwabl, 2004; although, again, many studies reported negative or no effect on the latter: von Engelhardt and Groothuis, 2011; Uller et al., 2005; Vergauwen et al., 2014). In contrast to males, we found no effect of experimental treatment on female reproductive traits at sexual maturity. Fewer studies have considered effects of yolk testosterone on female traits. In a large-scale field study, Ruuskanen et al. (2012a,b) found no long-term effect of yolk testosterone manipulation on any breeding parameter, in either sex, including laying date, clutch size, number of hatchlings and fledging success. Rubolini et al. (2007) reported that females hatching from testosterone-treated eggs had a lower egg-laying rate than controls, higher rates of egg infertility but no difference in egg size, and there was no sex difference in yolk size among the eggs laid by control females. In contrast, Uller et al. (2005) found that females experiencing relatively high levels of testosterone during embryonic development laid smaller eggs as adults.

The main goal of our study was to determine whether there were synergistic or antagonistic (compensatory) effects of multiple maternally derived egg components. We predicted synergistic effects of MAb and yolk testosterone for nestling and immature growth traits, and for adult 'quality' traits, but compensatory or even additive negative effects for immune function traits (Table 1). Synergy would be detected if offspring treated with both MAb and yolk testosterone had higher trait values than offspring treated with either MAb or yolk testosterone alone. We found no statistically significant evidence for this for any measured trait, and absolute trait values were only higher for one trait: 30 day mass in males (with the opposite pattern in females; see Fig. 2A). Compensatory effects of MAb and yolk testosterone should involve offspring treated with both MAb and yolk testosterone having intermediate trait values compared with offspring treated with either MAb or yolk testosterone alone. Although we found intermediate values for several traits, such as 26 day PHA response (see Fig. 3A) and brood size at fledging among manipulated clutches, none of these effects were significant. To our knowledge, there have only been two previous experimental studies that have attempted to simultaneously manipulate more than one egg component to look for interacting, synergistic or antagonistic effects of multiple egg traits. In Japanese quail, treatment of eggs with either exogenous carotenoids or testosterone had weak negative effects on offspring phenotype (e.g. lower hatching mass) but when eggs were treated simultaneously with both carotenoids and (th)-208t((th)-208tea
d (ma
ive pr (tra ion())-863.9 (an)18 (d)-122.3 (fo)16.3 uon
ma
(e)-1208
se(e)-11758.